## Appendix A Marked-up Version

Claims 1-5 and 18 were amended as follows:

- 1. (Amended) An isolated *dwf7* polynucleotide that imparts at least one *dwf7* mutant phenotype when expressed in a plant, said polynucleotide selected from the group consisting of (a) a polynucleotide comprising the nucleotide sequence depicted at positions 143 to 322, inclusive, of [Figures 8A-8D] <u>SEQ ID NO:20</u>; (b) a polynucleotide comprising the nucleotide sequence depicted at positions 143 to 1552, inclusive, of [Figures 8A-8D] <u>SEQ ID NO:20</u>; (c) a polynucleotide comprising a nucleotide sequence having at least 70% identity to the nucleotide sequence of (a) or (b); (d) a fragment of (a), (b) or (c) comprising at least 15 contiguous nucleotides; and (e) complements of (a), (b), (c), (d) or (e).
- 2. (Amended) The isolated *dwf7* polynucleotide of claim 1, wherein said polynucleotide comprises the nucleotide sequence depicted at positions 143 to 322, inclusive, of [Figures 8A-8D] <u>SEQ ID NO:20</u> or the complement thereof.
- 3. (Amended) The isolated *dwf7* polynucleotide of claim 1, wherein said polynucleotide comprises the nucleotide sequence depicted at positions 143 to 1552, inclusive, of [Figures 8A-8D] <u>SEQ ID NO:20</u> or the complement thereof.
- 4. (Amended) The isolated *dwf7* polynucleotide of claim 1, wherein said polynucleotide consists of the nucleotide sequence depicted at positions 143 to 322, inclusive, of [Figures 8A-8D] <u>SEQ ID NO:20</u> or the complement thereof.

- 5. (Amended) The isolated *dwf7* mutant polynucleotide of claim 1, wherein said polynucleotide consists of the nucleotide sequence depicted at positions 143 to 1552, inclusive, of [Figures 8A-8D] <u>SEQ ID NO:20</u> or the complement thereof.
- 18. (Amended) A method for altering the sterol composition of a plant relative to the wild-type plant comprising:
- (a) introducing [the] <u>a</u> polynucleotide [of claim 1] <u>comprising the nucleotide</u> sequence depicted at positions 143 to 322, inclusive, of SEQ ID NO:20 or a <u>polynucleotide comprising the nucleotide sequence depicted at positions 143 to 1552, inclusive, of SEQ ID NO:20</u> into a plant cell to produce a transformed plant cell; and
- (b) producing a transgenic plant from the transformed plant cell, said transgenic plant having an altered sterol composition relative to the wild-type plant.

New claims 36-51 have been added as follows:

- --36. (New) A recombinant vector comprising:
- (a) the isolated dwf7 polynucleotide of claim 2; and
- (b) control elements that are operably linked to said polynucleotide whereby a coding sequence within said polynucleotide can be transcribed and translated in a host cell, and at least one of said control elements is heterologous to said coding sequence.
  - 37. (New) A host cell transformed with the recombinant vector of claim 36.
  - 38. (New) A transgenic plant comprising the polynucleotide of claim 2.
  - 39. (New) A method of producing a transgenic plant comprising the steps of:

- (a) introducing the polynucleotide of claim 2 into a plant cell to produce a transformed plant cell; and
  - (b) producing a transgenic plant from the transformed plant cell.
  - 40. (New) A recombinant vector comprising:
  - (a) the isolated dwf7 polynucleotide of claim 3; and
- (b) control elements that are operably linked to said polynucleotide whereby a coding sequence within said polynucleotide can be transcribed and translated in a host cell, and at least one of said control elements is heterologous to said coding sequence.
  - 41. (New) A host cell transformed with the recombinant vector of claim 40.
  - 42. (New) A transgenic plant comprising the polynucleotide of claim 3.
  - 43. (New) A method of producing a transgenic plant comprising the steps of:
- (a) introducing the polynucleotide of claim 3 into a plant cell to produce a transformed plant cell; and
  - (b) producing a transgenic plant from the transformed plant cell.
  - 44. (New) A recombinant vector comprising:
  - (a) the isolated dwf7 polynucleotide of claim 4; and
- (b) control elements that are operably linked to said polynucleotide whereby a coding sequence within said polynucleotide can be transcribed and translated in a host cell, and at least one of said control elements is heterologous to said coding sequence.
  - 45. (New) A host cell transformed with the recombinant vector of claim 44.

- 46. (New) A transgenic plant comprising the polynucleotide of claim 4.
- 47. (New) A method of producing a transgenic plant comprising the steps of:
- (a) introducing the polynucleotide of claim 4 into a plant cell to produce a transformed plant cell; and
  - (b) producing a transgenic plant from the transformed plant cell.
  - 48. (New) A recombinant vector comprising:
  - (a) the isolated dwf7 polynucleotide of claim 5; and
- (b) control elements that are operably linked to said polynucleotide whereby a coding sequence within said polynucleotide can be transcribed and translated in a host cell, and at least one of said control elements is heterologous to said coding sequence.
  - 49. (New) A host cell transformed with the recombinant vector of claim 48.
  - 50. (New) A transgenic plant comprising the polynucleotide of claim 5.
  - 51. (New) A method of producing a transgenic plant comprising the steps of:
- (a) introducing the polynucleotide of claim 5 into a plant cell to produce a transformed plant cell; and
  - (b) producing a transgenic plant from the transformed plant cell.
- 52. (New) The method of claim 16, wherein the transgenic plant has a phenotype selected from the group consisting of altered cell length, altered periods of flowering, altered branching, altered seed production, altered leaf size, altered plant height, altered sterol composition, prolonged life, increased growth in the dark, GA modulation, and altered plant growth and development.

- 53. (New) The method of claim 52, wherein the phenotype is altered plant growth development is in the dark.
  - 54. (New) A method of producing a transgenic plant comprising the steps of:
- (a) introducing the polynucleotide having sequence ID No.19 into a plant cell to produce a transformed plant cell; and
  - (b) producing a transgenic plant from the transformed plant cell.
- 55. (New) The method of claim 54, wherein the transgenic plant has a phenotype selected from the group consisting of altered cell length, altered periods of flowering, altered branching, altered seed production, altered leaf size, altered plant height, altered sterol composition, prolonged life, increased growth in the dark, GA modulation, and altered plant growth and development.
- 56. (New) The method of claim 55, wherein the phenotype is altered plant growth development is in the dark. --